## REFERENCES

Those marked \* are reprints deposited at the Dorset County Museum Library.

- \*Bamber, R.N., Henderson, P.A. & Turnpenny, A.W.H. (1985) The early life history of the sand smelt (<u>Atherina presbyter</u>). J. mar. biol. Ass. U.K., 65: 697-706.
- \*Bamber, R.N. & Henderson, P.A. (1985a) Morphological variation in British Atherinids and the status of <u>Atherina presbyter</u> Cuvier (Pisces: Atherinidae). Biological Journal of the Linnean Society, 25: 61-76.
- \*Bamber, R.N. & Henderson, P.A. (1985b) Diplostomiasis in sand smelt, <u>Atherina</u> <u>presbyter</u> Cuvier, from the Fleet, Dorset and its use as a population indicator. J. Fish Biol., 26: 223-229.
- \*Bass, J.A.B., Beaumont, W.R.C., Ladle, M. & Mann, R.H.K. (1983) Investigation of the Fleet, Weymouth, as a potential spawning and nursery area for the bass, <u>Dicentrarchus</u> <u>labrax</u> (L.). Report commissioned by Dorset County Council, 19pp.
- \*Bird, E.C.F. (1972) The Physiography of the Fleet. Proc. Dorset nat. Hist. archaeol. Soc., 93: 125-131.
- \*Burrows, E.M. (1981) The algae of the Fleet. In: The Fleet and Chesil Beach (ed. M. Ladle), Dorset County Council.
- \*Butcher, R.W. (1933a) Notes on the variation of the British species of Zostera. Report of the Botanical Exchange Club.
- \*Butcher, R.W. (1933b) Studies on the ecology of rivers. No. 1. On the distribution of macrophytic vegetation in the rivers of Britain. J. Ecol., 21.
- \*Butcher, R.W. (1941a) Zostera Report on the Present (Feb. 1933) condition of Eel Grass in the coats of England (based on a survey during Aug-Oct 1933). International Wildfowl Inquiry 1941.
- \*Butcher, R.W. (1941b) The distribution of <u>Zostera</u> (Eel Grass, Widgeon Grass) and other seashore plants in relation to the migration of wildfowl. International Wildfowl Inquiry 1941.
- \*Brandwood, A. (1982) Intraspecific variation and distribution of <u>Littorina</u> <u>rudis</u> (Maton) in the Fleet - a coastal lagoon in Dorset. Proc. Dorset nat. Hist. archaeol. Soc., 104: 165-167.
- \*Brandwood, A. (1985) The effects of environment upon shell construction and strength in the Rough periwinkle <u>Littorina</u> <u>rudis</u> Maton (Mollusca:Gastropoda). J. Zool. Lond. (A), 206: 551-565.
- \*Campbell, D. (1985) The life cycle of <u>Gymnophallus</u> <u>rebecqui</u> (Digenea:Gymnophallidae) and the response of the bivalve <u>Abra</u> <u>tenuis</u> to its Metacercariae. J. mar. biol. Ass. U.K., 65: 589-601.
- \*Carr, A.R. (1969) Size grading along a pebble beach, Chesil Beach, England. J. Sedimentary Petrology, 29, No. 1, March 69.

- \*Carr, A.R. (1971) Experiments on longshore transport and sorting of pebbles, Chesil Beach, England. J. Sedimentary Petrology, 41, No. 4, Dec. 1971.
- Carr, A.P. & Blackey, M.W.L. (1973). Ideas on the origin and development of Chesil Beach, Dorset. Proc. Dorset nat. Hist. archaeol. Soc., 95: 9-17.
- \*Carr, A.R. (1983) Scientific conservation as a minority interest. The example of Chesil Beach. Earth Sc. Conservation No. 2 1983.
- \*Carr, A.R., Gleason, R. & King, A. (1970) Significance of pebble size and shape in sorting by waves. Sedimentary Geology, 194909: 89-101.
- Claridge, P.M. & Potter, I.C. (1983) Movements, abundance, age composition and growth of bass, <u>Dicentrarchus</u> <u>labrax</u>, in the Severn estuary and inner Bristol Channel. J. mar. biol. Ass. U.K., 63: 871-879.
- Critchley, A.T., Farnham, W.F. & Morrell, S.L. (1983) A chronology of new European sites of attachment for the invasive brown alga, <u>Sargassum</u> muticum, 1973-1981. J. mar. biol. Ass. U.K., 63: 799-811.
- Demestre, M., Roig, A., Sostoa, A. de & Sostoa, F.J. de (1977) Contribucio al'edtudi de la ictiofauna continental del delta de l'Ebre. Treb. Inst. Cat. Hist. Nat., 8: 145-226.
- Dyrynda, P.E.J. (1984) Investigation of the subtidal ecology of the Fleet Lagoon (Dorset). Report to the Nature Conservancy Council.
- Dyrynda, P.E.J. (1985) Poole Harbour subtidal survey Southern Sector -1984. Report to the Nature Conservancy Council.
- \*Dyrynda, P.E.J. & Farnham, W.F. (1984) Benthic communities of a rapids system within the Fleet lagoon, Dorset. Progress in Underwater Sci., 10: 65-82.
- \*Farnham, W. (1985) "Japanese Seaweed" <u>Sargassum muticum</u>. Interim report to MSC.
- \*Farnham, W., Blunden, G. & Gorden, S.M. (1985) Occurrence and pigment analyses of the sponge endobiont <u>Microspora</u> <u>ficulina</u> Chlorophyceae. Bot. marina, 17: 799-81.
- \*Fitzpatrick, V.M. (1976) The Fleet Study Group's 1st Progress Report.
- Forbes, E. & Hanley, S. (1848-53). A History of British Mollusca and their Shells. London, Van Voost.
- \*Gibbs, P.E. (1982) The turbellarian <u>Paravortex</u> parasitic in <u>Scrobicularia</u> bivalves: new records from S.W. England. J. mar. biol. Ass. U.K., 62: 739-740.
- \*Gibbs, P.E. (1984) The population cycle of the bivalve <u>Abra tenuis</u> and its mode of reproduction. J. mar. biol. Ass. U.K., 64: 791-800.
- \*Gleason, R. & Hardcastle, P.V. (1973) The significance of wave parameters in the sorting of beach pebbles. Estuarine and Coastal Science, 1: 11-18.
- Gray, J.S. & Rieger, R.M. (1971) A quantitative study of the meiofauna of an exposed sandy beach at Robin Hood's Bay, Yorkshire. J. mar. biol. Ass. U.K., 51: 1-19.

- Guiart, J. 1901. Contribution a l'etude des gasteropodes opisthobranches et en particulier des cephalaspides. Mem. Soc. zool. France, 14: 5-219.
- \*Hardcastle, P.V. & King, A.C. (1972) Chesil Beach sea wave records. Civil Engineering, March 1972.
- Harris, R.P. (1972) The distribution and ecology of the interstitial meiofauna of a sandy beach at Whitsand Bay, East Cornwall. J. mar. biol. Ass. U.K., 51: 1-18.
- \*Hibbard, D.V. (1977) Observations on the ultrastructure of the crypotmonad endosymbiont of the red water ciliate <u>Mesodinium</u> rubra. J. mar. biol. Ass. U.K., 57: 45-61.
- \*Holme, N.A. & Bishop, G.M. (1980) Report on the sediment shores of Dorset, Hampshire and the Isle of Wight. SMBA/MBA Intertidal Survey Unit, Jan. 1980.
- \*Holmes, N.T.H. (1983) The distribution of <u>Zostera</u> and <u>Ruppia</u> in the Fleet. Report to NCC.
- \*Holmes, N.T.H. (1984) Chesil Beach and the Fleet. The Living Countryside, No. 164. 1984.
- \*Humphrey, E.C. (1981) A preliminary survey of the meiofauna of the Fleet. MSc Dissertation, Dept of Oceanography, Southampton University, 91pp.
- \*Ilchester, The Earl of (1934) The Abbotsbury Swannery. Proc. Dorset nat. Hist. archaeol. Soc., Feb. 1934.
- \*Jackson, S.P. (1924) A report on the Tern colony of the Chesil Beach for DNT.
- Kelley, D.F. (1979) Bass populations and movements on the west coast of the U.K. J. mar. biol. Ass. U.K., 59: 889-936.
- Kennedy, M. & Fitzmaurice, P. (1972) The biology of the bass, <u>Dicentrarchus</u> <u>labrax</u>, in Irish waters. J. mar. biol. Ass. U.K., 52: 557-597.
- Kitching, J.A. & Ebling, F.J. (1967) Ecological studies at Lough Ine. Advances in Ecological Research, 4: 197-291.
- \*Ladle, M. (1981a) Structure and biology of a unique coastal feature. In: The Fleet and Chesil Beach (ed. M. Ladle), Dorset County Council.
- \*Ladle, M. (1981b) Invertebrate fauna of the Fleet. In: The Fleet and Chesil Beach (ed. M. Ladle), Dorset County Council.
- Lewis, J.R. & Powell, H.T. (1960) Aspects of the intertidal ecology of rocky shores in Argyll, Scotland: I and II. Trans. Roy. Soc. Edinb., 64: 7-74.
- Mare, M.F. (1942) A study of a marine benthic community with special reference to the micro-organisms. J. mar. biol. Ass. U.K., 25: 519-554.
- McIntyre, A.D. & Murison, D.J. (1973) The meiofauna of a flatfish ground. J. mar. biol. Ass. U.K., 53: 93-118.
- \*Moore, J.A. (1979) The current status of the Characeae (stoneworts) in the British Isles. Watsonia, 12: 297-309.

- Morton, J.E. (1972) The form and functioning of the pallial organs in the opisthobranch <u>Akera bullata</u> with a discussion on the nature of the gill in Notaspidea and other tectibranchs. The Veliger, 14: 337-349.
- Morton, J.E. & Holme, N.A. (1955) The occurrence at Plymouth of the opisthobranch <u>Akera bullata</u>, with notes on its habits and relationships. J. mar. biol. Ass. U.K., 34: 101-112.
- Nikouyan, A. (1980) Factors affecting the microdistribution of meiofauna on a sandy shore. M.Sc. Dissertation, University of Southampton, 74pp.
- \*Pickering, C. (in press) A survey of the distribution and community structure of the marine benthic invertebrates of the Fleet - brackish coastal lagoon in Dorset. Proc. Dorset nat. Hist. archaeol. Soc.
- \*Robinson, I.S. (1981a) Tides and water levels in the Fleet. In: The Fleet and Chesil Beach (ed. M. Ladle), Dorset County Council.
- \*Robinson, I.S. (1981b) Salinity structure and tidal flushing of the Fleet. In: The Fleet and Chesil Beach (ed. M. Ladle), Dorset County Council.
- Robinson, I.S. (1983) A tidal flushing model of the Fleet an English tidal lagoon. Estuarine, Coastal and Shelf Science, 16: 669-688.
- Robinson, I.S., Warren, L. & Longbottom, . . (1983) Sea-level fluctuations in the Fleet, an English tidal lagoon. Estuarine, Coastal and Shelf Science, 16: 651-668.
- Russell, F.S. (1976) The eggs and planktonic stages of British marine fishes. Academic Press, London. 324pp.
- Russell, F.S. (1980) On the distribution of postlarval fish in the Bristol Channel. Bull. mar. Ecol., 8: 283-290.
- \*Seaward, D.R. (1978) The Marine Molluscs of the Fleet, Dorset. Proc. Dorset nat. Hist. archaeol. Soc., 100: 100-108.
- Seaward, D.R. (1979) <u>Alderia modesta</u> on a sandbank at L.W.S.T. The Conchologists Newsletter No. 69: 155-156.
- Seaward, D.R. (1982) Sea Area Atlas of the Marine Molluscs of Britain and Ireland. Published for the Conchological Society of Great Britain and Ireland by the Nature Conservancy Council.
- \*Seaward, D.R. (1983) The distribution of Marine Molluscs in the Fleet, Dorset. Report to NCC. Oct. 1983.
- Seaward, D.R. (1985) <u>Emarginula rosea</u> and <u>Lepidopleurus asellus</u> living intertidally in Dorset, with a note on <u>Truncatella</u> <u>subcylindrica</u>. The Conchologists' Newsletter No. 95: 322-323.
- \*Smith, G., Moryson, C.J. & Walmsley (1975) <u>Clostridium botulinum</u> in the lakes and waterways of London. J. Hyg. Camb., 75: 371.
- \*Smith, G.R., Moryson, C.J. & Walmsley (1977a) The low prevalence of <u>C</u>. <u>botulinum</u> in the lakes, marshes and waterways of the Camargue. J. Hyg. Camb., 78: 33.

- \*Smith, G.R., Moryson, C.J. & Walmsley (1977b) A comparison of the distribution of <u>C</u>. <u>botulin um</u> in soil and in lake mud. J. Hyg. Camb., 78: 39.
- \*Smith, G.R., Moryson, C.J. & Walmsley (1977c) Avian pollution and the high prevalence of <u>C. botulin um</u> in the Norfolk Broads. The Vet Record, 100(6): 106-109, Feb. 5.
- \*Smith, G.R., Moryson, C.J. & Walmsley (1977d) <u>Clostridium botulin um</u> type D in Britain. Vet. Record, Feb. 5.
- \*Smith, G.R., Moryson, C.J. & Walmsley (1978) <u>C. botulin um</u> in aquatic environments in Great Britain and Ireland. J. Hyg. Camb., 80: 431.
- \*Stephens, J.M. (1969) The distribution of Recent Foraminifera in the Fleet, Dorset. Unpublished M.Sc. Thesis, University of Wales (Aberystwyth).
- \*Sturdy, R. (1972) The Fleet Survey an investigation of conservation and recreational issues affecting the Fleet and surrounding area. Aug. 1972.
  - Thompson, T.E. (1976) Biology of Opisthobranch Molluscs I. London, Ray Society.
- \*Tittley, I., Farnham, W., Fletcher, R.L., Morrell, S. & Bishop, G. (1985) Sublittoral seaweed assemblages of some northern Atlantic Islands. Progress Underwater Sci., 10: 39-52.
- \*Vaughan, T. (1984) Bridge over shifted waters. Heavy Construction Nov./Dec. 1983.
- \*Whittaker, J.E. (1968) CEGB's data 1968 survey for possible Nuclear Power site. Photocopies for J. Whittaker.
- \*Whittaker, J.E. (1969) 2 sets maps and figures from Whittaker's thesis 1969.
- \*Whittaker, J.E. (1972) Recent brackish and marine Ostracoda from the Fleet and Weymouth Bay, Dorset. Ph.D Thesis.
- \*Whittaker, J.E. (1973) 9 aerial photographs done by RAF.
- \*Whittaker, J.E. (1978) 2 copies, The Fleet, Dorset a seasonal study of the watermass and its vegetation. Proc. Dorset. nat. Hist. archaeol. Soc., 100: 73-99.
- \*Whittaker, J.E. (1981a) The hydrology of the Fleet. In: The Fleet and Chesil Beach (ed. M. Ladle), Dorset County Council.
- \*Whittaker, J.E. (1981b) The distribution of <u>Zostera</u> and <u>Ruppia</u> in the Fleet. In: The Fleet and Chesil Beach (ed. M. Ladle), Dorset County Council.
- \*Whittaker, J.E. (1981c) The distribution of ostracoda in the Fleet. In: The Fleet and Chesil Beach (ed. M. Ladle), Dorset County Council.
- \*Whittaker, J.E. & Farnham, W.F. (1983) The structure and function of brackish water and inshore communities. EBSA Heriot-Watt Symposium, Edinburgh 1985, paper no. 5.